

RESEARCH ARTICLE

Open Access

Perceived Stress, Sources and Severity of Stress among medical undergraduates in a Pakistani Medical School

Mohsin Shah^{1*}, Shahid Hasan², Samina Malik², Chandrashekhar T Sreeramareddy³

Abstract

Background: Recently there is a growing concern about stress during undergraduate medical training. However, studies about the same are lacking from Pakistani medical schools. The objectives of our study were to assess perceived stress, sources of stress and their severity and to assess the determinants of stressed cases.

Methods: A cross-sectional, questionnaire-based survey was carried out among undergraduate medical students of CMH Lahore Medical College, Pakistan during January to March 2009. Perceived stress was assessed using the perceived stress scale. A 33-item questionnaire was used to assess sources of stress and their severity.

Results: The overall response rate was 80.5% (161 out of 200 students). The overall mean perceived stress was 30.84 (SD = 7.01) and was significantly higher among female students. By logistic regression analysis, stressed cases were associated with occurrence of psychosocial (OR 5.01, 95% CI 2.44-10.29) and academic related stressors (OR 3.17 95% CI 1.52-6.68). The most common sources of stress were related to academic and psychosocial concerns. 'High parental expectations', 'frequency of examinations', 'vastness of academic curriculum', 'sleeping difficulties', 'worrying about the future', 'loneliness', 'becoming a doctor', 'performance in periodic examinations' were the most frequently and severely occurring sources of stress. There was a negative but insignificant correlation between perceived stress and academic performance ($r = -0.099$, $p > 0.05$).

Conclusion: A higher level of perceived stress was reported by the students. The main stressors were related to academic and psychosocial domains. Further studies are required to test the association between stressed cases and gender, academic stressors and psychosocial stressors.

Background

Recently stress during medical training is increasingly being reported in published literature. Previous studies have shown fairly high levels of distress, such as symptoms of depression [1,2] and even suicide thoughts [3,4] among medical undergraduates. The potential negative effects of emotional distress on medical students include impairment of functioning in class-room performance and clinical practice, stress-induced disorders and deteriorating performance [5,6]. Perceived medical stress has also been linked to current mental distress [7] and to forthcoming health problems [8]. However, there is very

little information about the effect of stress on academic performance during medical training [9].

In addition to stress the students' social, emotional and physical as well as family problems may influence their learning ability and academic performance [10,11]. Higher levels of stress may have a negative impact on the students learning ability. Excessive stress may result in mental and physical problems and may diminish a student's sense of worth and might affect his/her academic achievement [12,13]. Earlier studies have classified stressors into three main categories: academic pressures, social issues and financial problems [14]. Studies from developing countries like Thailand, India, Malaysia and even Pakistan have reported stress among medical students and have also underscored the role of academics as a potential stressor [15-18]. But these studies either did not use the perceived stress scale or did

* Correspondence: mohsin255@hotmail.com

¹3rd year Undergraduate Medical Student, CMH Lahore Medical College, University of Health Sciences, Abdur Rahman Road, Lahore Cantt, Pakistan

not study the correlation between stress and academic performance [19]. Studies have reported that academic constraints, [20] and also factors like age, gender, ethnicity and marital status may also influence students' severity of stress and hence academic performance [21,22].

At the Combined Military Hospital Medical College (CMH LMC), Lahore, Pakistan, there are students from Pakistan, Saudi Arabia, United States, U.A.E and other countries. These students come from diverse cultural, socioeconomic and educational backgrounds. These foreign nationals who account for more than a third of the student population are exposed to a new learning environment, making new social circles and also adapting to a new and different world during their training at CMH LMC. This may be a very stressful experience especially during the formative first and second years of their course. The relative paucity of information about stress and its sources during the early years of medical undergraduate training in Pakistan warranted this study. The objectives of our study were:

1. To assess perceived stress, sources of stress and their severity.
2. To assess the determinants of stressed cases.

Methods

Setting and participants

The present study was undertaken at CMH LMC, affiliated to University of Health Sciences (UHS), Lahore. The three basic science subjects (Anatomy, Physiology and Biochemistry) are taught during first two years of MBBS (Bachelor of Medicine and Bachelor of Surgery) course in a lecture-based region wise manner. Each year CMH LMC admits a batch of 100 students for the MBBS course. At the end of each academic year the students sit for their professional examinations, held by the UHS, to qualify for the next academic year. The students of first and the second year MBBS admitted during 2006 and 2007 respectively were invited to participate in this study

Study Design

A cross-sectional survey using self-administered questionnaire. The institutional ethics committee of CMH LMC approved this study. A total of 200 medical undergraduates from the first and second year MBBS batches at CMH Lahore Medical College were included for survey. We did not have any earlier estimates of prevalence of stress among medical students in Pakistan. Therefore we assumed the prevalence to be 50% and calculated the required sample for our survey with an allowable error of 15%, 95% confidence limits and 10% for non-responders. Thus the final sample size required was 187. The students were asked to complete a set of

questionnaires consisting of three parts namely: demographic information and academics, (Appendix 3) perceived stress scale [19] (Appendix 4), a 33-item list of potential stressors (Appendix 5).

Definitions of variables

Perceived stress

Perceived stress was measured using the perceived stress scale (PSS-14) [19], which comprised of 14 questions with responses varying from 0 to 4 for each item and ranging from never, almost never, sometimes, fairly often and very often respectively on the basis of their occurrence during one month prior to the survey. The PSS has an internal consistency of 0.85 (Cronbach α co-efficient) and test-retest reliability during a short retest interval (several days) of 0.85 [19]. It assesses the degree to which participants evaluate their lives as being stressful during the past month. It does not tie appraisal to a particular situation; the scale is sensitive to the nonoccurrence of events as well as ongoing life circumstances. PSS-14 scores are obtained by reversing the scores on four positive items, for example 0 = 4, 1 = 3, 2 = 2, etc. and then summing across all 14 items. Items 4, 5, 6, 7 and 10 are the positively stated items. The scale yielded a single score with high scores indicating higher levels of stress and lower levels indicating lower levels of stress. The PSS-14 has a possible range of scores from 0 to 56. The range of PSS scores were also divided into stratified quartiles. The upper two and lower two quartiles were combined (28 being the operational cut off value for the upper bound) and were labeled as stressed and not stressed respectively. This cut off value was selected in accordance to a similar study from Egypt [23].

Academic performance

Academic performance was measured using the student's self reported UHS professional examination score. The total university examination score (marks) was out of 550. The examination score was measured using an open ended question and the possible range of scores could have been from 0 to 550.

Stressors

The potential stressors in the questionnaire were adapted from a similar study from Nepal by Sreeramreddy et al [24]. A total of 33 stressors were listed and were grouped as academic, psychosocial and health related. For each potential stressor the frequency of occurrence was classified as never, rarely, sometimes, often and always and scored as 1,2,3,4 and 5 respectively. The severity of each stressor was rated using a Likert scale (1-10) ranging from not severe to very severe. The students were asked to indicate if any of the stressors had been affecting them.

Data collection

One hundred students each from first and second year MBBS admitted during 2006 and 2007 respectively were invited to participate in this survey. The students were given a copy of the written instructions and objectives of the study (Appendix 1). Informed consent was taken from all the participants (Appendix 2). The participants were assured of confidentiality of the information provided and had an option of refusal to participate in the survey. The anonymous questionnaire was distributed amongst students during breaks from their teaching schedule and the researchers collected the completed questionnaires.

Data analysis

The data was analysed using Statistical Package for Social Sciences (SPSS) 17.0 for Windows (SPSS, Inc., Chicago, IL, USA). The mean score of perceived stress were calculated. The number and percentage of stressed cases were calculated according to demographic variables. We grouped the frequency of occurrence of stressors as never/rarely, sometimes, and often/always. Percentage frequency of occurrence was calculated for each of the stressors from academic, psychosocial and health domains. Descriptive statistics were calculated for severity of stressors. Pearson product-moment correlation was applied to test the correlation between perceived stress (PSS score) and self-reported academic performance. Logistic regression analyses were carried out to assess determinants of stressed cases. We considered perceived stress (stressed cases) as the dependent variable, demographic variables and groups of stressors (i.e. academic, psychosocial and health-related) as the independent variables. Adjusted odds ratios (OR), 95% confidence intervals (95% CI) were calculated. A p -value < 0.05 was considered as significant.

Results

Demographic characteristics of the respondents

Out of 200 students 161 completed and returned the questionnaire giving an overall response rate of 80.5% (78% for first year and 83% for second year). The mean age was 20.35(SD = 1.09) with a range of 17-27 years. Fifty three students were male (32.92%) and 108 were females (67.08%). Among the nationalities 121(75.16%) were Pakistani and 40 (24.8%) were foreign nationals [Saudi Arabia (13), USA (11), U.A.E (6), Bahrain (3), Oman (4), Canada (1), Kuwait (1) and Qatar (1)]. Among the Higher Secondary School Certification 114 had completed their Fellow of science (Fsc) (69.57%), 40 had done A-levels (24.84%) while nine had completed their High School Diploma (5.59%). 148 students had gone to English medium schools (91.93%) whereas only 13 had attended Urdu (a local vernacular language) medium school (8.07%). Ninety nine students were day

scholars (61.49%) while 62 (38.51%) were residing in the hostels (Boarders). One hundred and thirty two (81.99%) students reported as being single (81.99%), 29 (18.01%) were involved in active relationships while none were married.

Perceived stress

Mean PSS score in the study population was 30.84(SD = 7.01) with a median of 31.00 (IQR 26-36). Mean PSS score for female students ($n = 108$) was 31.94 (SD 6.28) while the same for male students ($n = 53$) was 28.60 (SD 7.92). The 14-question survey instrument's sample response frequencies are given (Table 1). Using univariate analysis, only gender was significant ($p < 0.05$) with PSS score; the female students reported significantly higher levels of perceived stress than their male counterparts.

Academic performance

The mean academic score of the students was 381 (SD = 42.26). The mean academic score of the female students ($n = 108$) was 378.57 (SD 44.22); that of the male students ($n = 53$) was 387.15 (SD 37.71).

Sources and self-rated severity of stressors

The most frequently [number of respondents (percentages)] occurring sources of stress reported by students as often/always were, 'high parental expectations' [101 (63%)], 'frequency of examinations' [95(59%)], 'vastness of academic curriculum' [81(50%)], 'sleeping difficulties' [77 (48%)], 'performance in periodic examinations' [74 (46%)], 'worrying about the future' [72(45%)], 'loneliness' [66(41%)], 'becoming a doctor (expectations on all fronts)' [64(40%)], 'Non availability of learning materials' [56(35%)], 'quality of food in mess' [56(35%)]. 'High parental expectations', 'frequency of examinations', 'sleeping difficulties', 'dissatisfaction with the class lectures', 'worrying about the future', 'vastness of academic curriculum/syllabus', and 'loneliness' were rated as most severe (Table 2).

The correlation between perceived stress and academic performance was assessed using Pearson correlation coefficient; this was negative and not significant ($r = -0.099$, $p > 0.05$).

Determinants of stressed-cases by logistic regression

By logistic regression analysis stressed cases were associated with being a female [OR 2.25, 95% CI 1.13-4.49], occurrence of psychosocial [OR 5.01 (95%CI 2.44-10.29)] and academic-related stressors [OR 3.17 (95%CI 1.52-6.68)] (Table 3).

Discussion

In our study, we evaluated perceived stress among medical students including its sources and severity along with its correlation with academic performance, which may be of importance to both medical teachers and psychologists. To our knowledge such a detailed study has not

Table 1 Medicine students' responses to the perceived stress scale

Statement	Never	Almost never	Sometimes	Often	Very often
In the last month, how often have you been upset because of something that happened unexpectedly?	8 (5)	19 (12)	60 (37)	40 (25)	34 (21)
In the last month, how often have you felt that you were unable to control the important things in your life?	17 (11)	26 (16)	55 (34)	36 (22)	27 (17)
In the last month, how often have you felt nervous and "stressed"?	3 (2)	6 (4)	44 (27)	59 (37)	49 (30)
In the last month, how often have you dealt successfully with day to day problems and annoyances?	29(18)	50 (31)	59 (37)	21 (13)	2 (1)
In the last month, how often have you felt that you were effectively coping with important changes that were occurring in your life?	24 (15)	47 (29)	54 (34)	29 (18)	7 (4)
In the last month, how often have you felt confident about your ability to handle your personal problems?	36 (22)	47 (29)	53 (33)	18(11)	7 (4)
In the last month, how often have you felt that things were going your way?	15 (9)	38 (24)	70 (43)	25 (16)	13 (8)
In the last month, how often have you found that you could not cope with all the things that you had to do?	10 (5)	15 (9)	69 (43)	43 (27)	24 (15)
In the last month, how often have you been able to control irritations in your life?	5 (3)	21 (13)	73 (45)	39 (24)	23 (14)
In the last month, how often have you felt that you were on top of things?	8 (5)	22 (14)	57 (35)	59 (37)	15 (9)
In the last month, how often have you been angered because of things that happened that were outside of your control?	9 (6)	16 (10)	48 (30)	47 (29)	41 (25)
In the last month, how often have you found yourself thinking about things that you have to accomplish?	0 (0)	8 (5)	23 (14)	38 (24)	92 (57)
In the last month, how often have you been able to control the way you spend your time?	33 (20)	36 (22)	49 (30)	33 (20)	10 (6)
In the last month, how often have you felt difficulties were piling up so high that you could not overcome them?	15 (9)	31 (19)	57 (35)	27 (17)	31 (19)

Figures in each cell depict number of responses and their percentage for each question in PSS

been reported from Pakistani medical schools. In our study, medical students reported a higher level of perceived stress, which was significantly higher among female students, but it's correlation with self-reported academic performance was not significant. Academic and psychosocial stressors were reportedly more common both in terms of frequency of occurrence and severity. By logistic regression analyses stressed cases were associated with female gender, psychosocial and academic related stressors.

Life of a medical student or a health care professional can be very stressful. Mild, moderate, and high levels of stress and even burnout have been reported amongst medical students and health care professionals from other countries [25-29]. Previous studies have shown that medical school stress is a good predictor of nervous symptoms even when psychosocial variables such as marital status or cohabitation status, confident and other general self-esteem are taken into consideration [6]. Apart from all this, end-of-the-year first year medical students have shown to be worse off psychosocially than they were when they entered [30].

The amount and severity of stress experienced by medical students may vary according the settings of the medical school, the curricula, evaluation (examination) system etc. Previous studies from medical schools in different countries have reported varying levels of stress [1,5,6,9,13,15,16,18]. A study from Agha Khan

University, Pakistan has reported that more than 90% of its students experienced stressed at one time or the other during their course [18]. A similar study from India reported that 73% of the students had perceived stress at some point or the other during their medical schooling [15]. Saipanish reported that 61.4% of students in a Thai Medical School had come across some degree of stress as calculated by the Thai Stress Test [16]. These studies have used different instruments to measure stress. This limits the comparability among these studies. We chose the perceived stress scale since this instrument has been documented for its reliability and validity [19,31-42]. The advantage of PSS is that it can be applied to a wide range of settings, to different subject types and includes items measuring reactions to stressful situations as well as measures of stress [19,31-42]. An important limitation of other reviewed stress scales for health professions students is that it focuses only on academic stressors, and lack of inclusion of personal issues or reactions to stressful situations (psychosocial issues), and poor applicability to broader settings.

In our study, sample proportion (67.08%) of the female students was higher than their male (32.92%) counterparts. Mean PSS scores among female students was significantly higher than that of male students. A plausible reason may be the conservative nature of society in Pakistan, where women lack freedom to

Table 2 Response pattern of the 33 sources of stress and perceived severity (rated in a likert scale of 1-10) as reported by the students

Sources of Stress	Frequency of occurrence			Severity	
	Never/Rarely (%)	Sometimes (%)	Often/Always (%)	Median	IQR*
Academic Stressors					
Frequency of Examinations		20	21	59	6 3 - 8
Performance in Examinations		18	36	46	6 3 - 7
Academic Curriculum		20	30	50	5 3 - 7
Dissatisfaction with Class Lectures		29	42	29	5 3 - 7
Non-Availability of Adequate learning materials		39	26	35	5 2 - 7
Becoming a Doctor		30	30	40	5 2 - 8
Lack of time for recreation		36	29	35	5 2 - 7
Competition with Peers		43	23	34	4 1 - 7
Performance in practicals		65	16	19	2 1 - 4
Lack of special guidance from faculty		68	20	12	2 1 - 4
Psychosocial Stressors					
High Parental Expectations		24	13	63	7 3 - 10
Loneliness		30	29	41	5 2 - 8
Family Problems		46	29	25	4 1 - 7
Accommodation away from home		41	24	35	4 1 - 8
Political situation in the country		47	21	32	4 2 - 7
Relations with the Opposite Sex		55	21	24	3 1 - 7
Difficulty reading text books		63	26	11	3 1 - 5
Lack of entertainment in the institution and Lahore itself		52	24	24	3 1 - 6
Difficulty in the journey back home		49	17	34	3 1 - 7
Quality of food in mess		44	21	35	3 1 - 8
Financial strain		53	24	23	3 1 - 6
Inability to socialize with peers		63	20	17	3 1 - 5
Living conditions in the hostel		63	15	22	1 1 - 5
Member of fraternity or sorority		83	6	11	1 1 - 2
Lack of personal interest in medicine		76	14	10	1 1 - 3
Adjustment with roommate/s		69	10	21	1 1 - 5
Health Related Stressors					
Sleeping Difficulties		27	25	48	6 3 - 8
Class Attendance		49	23	28	4 1 - 6
Nutrition		43	25	32	4 2 - 7
Exercise		46	22	32	4 1 - 7
Quality of food in mess		44	21	35	3 1 - 8
Physical disability		63	21	16	2 1 - 5
Alcohol/Drug abuse/Smoking		86	2	12	1 1 - 1

*Inter Quartile Range

Table 3 Determinants of stressed cases by logistic regression

Determinants	Number	Number of stressed Cases (%)	Univariate OR (95% CI)	Adjusted OR* (95% CI)
Age (in completed years)				
< 21	95	67(71.2)	1	1
>=21	66	41(62.1)	0.69 (0.33-1.4)	0.69 (0.35-1.33)
Gender				
Male	53	29(54.7)	1	1
Female	108	79(73.1)	2.25 (1.07-4.76)	2.25 (1.13-4.49)
Batch of Study				
First	83	55(66.3)	1	1
Second	78	53(67.9)	1.08 (0.53-2.20)	1.08 (0.56-2.08)
Nationality				
Pakistan	121	81(66.9)	1	1
International	40	27(67.5)	1.03 (0.45-2.36)	1.03 (0.48-2.19)
Marital Status				
Single	132	85(64.4)	1	1
In a relationship	29	23(79.3)	2.12 (0.75-6.28)	2.12 (0.81-5.57)
Medium of teaching in School				
English	148	98(66.2)	1	1
Urdu	13	10(76.9)	1.70 (0.41-8.19)	1.70 (0.45-6.46)
HSSC				
A-levels	40	30(75)	1	1
Fsc and HSD	121	78(64.4)	0.21 (0.09-0.49)	0.62 (0.28-1.41)
Day Scholar/Boarder				
Day Scholar	99	68(68.7)	1	1
Boarder	62	40(64.5)	0.83 (0.40-1.71)	0.83 (0.42-1.62)
Occurrence of Psychosocial Stressors				
Less than often	50	19(38.0)	1	1
Often/Always	111	87(78.3)	5.91 (2.69-13.13)	5.01 (2.44-10.29)
Occurrence of Academic Stressors				
Less than often	97	54(55.7)	1	1
Often/Always	64	52(81.3)	3.45 (1.55-7.8)	3.17 (1.51-6.68)
Occurrence of Health Stressors				
Less than often	147	97(65.9)	1	1
Often/Always	14	9(64.2)	0.93 (0.26-3.39)	0.87 (0.28-2.75)

*Adjusted with Age, Gender, Batch of Study, Nationality, Marital Status, Medium of teaching in school, HSSC, Day Scholar/Boarder, Occurrence of Psychosocial Stressors, Occurrence of Academic Stressors, Occurrence of Health Stressors

participate in extra-curricular activities owing to the restrictions imposed on them by the society. However, Cohen has reported that there was no significant difference in stress using PSS between male and female students. In addition, private life may conflict with professional life and this may cause stress. We lacked sufficient information, which could assist us in carrying out further analysis about this. The overall average stress score in our sample was significantly higher ($p < 0.05$) than that reported by Cohen [31].

In our study the correlation between perceived stress and academic performance was negative and not statistically significant. This finding is contrary to existing literature which states that acute stress is a predictor of reduced scholastic performance, especially in

examinations [9,43]. Plausible explanations for such contradictory reports could be that only acute stress but not longstanding stress may affect academic grades during examinations. In addition, the students who are striving to perform well in examinations may make themselves stressed. Individual coping styles and skills along with their access to different forms of social support may have played a role in negating the effect of stress on academic performance; this could not be explored in our study.

Most students had experienced either academic and/or psychosocial stressors. Among academic stressors, 'tests/exams' were the chief sources of stress. Despite this, test/exams are important in the medical training as a standard for evaluation/assessment. Also examinations encourage students' learning and also provide feedback to the

teachers. Those students who perceive tests/exams as a burden may experience stressful situations while for others, who consider exams useful, they may assist in their learning. Previous studies have also reported that academics/exams are common sources of stress among medical students [9,15-19,44-47]. There may be a need to revisit the evaluation/examination system to make it less stressful to the students. An important source of stress was related to psychosocial factors. This may be due to time constraints for self, family, friends and entertainment due to the demanding medical curriculum. Another reason could be inadequate recreational facilities provided by the college as reported by the students. The students, who were stressed, reported that psychosocial and academic related stressor groups had occurred more frequently. This was revealed by the association of these factors with stressed cases by logistic regression analysis. This may suggest that students had a global response to a wide range of potential stressors rather than being limited to a few specific items. However, there was a high odds ratio on both univariate and multivariate analysis suggesting high collinearity. This suggests that students having academic stressors were more likely to have psychosocial stressors as well or vice versa. A study from USA has recommended that teaching stress management and self-care skills to medical students may prove to be beneficial [48]. There is a need to look at the applicability of such measures which are feasible in our medical school setting.

Limitations

Lack of generalisation of our results to other medical schools in Pakistan is an important limitation of this study. Since the information was collected on self-administered questionnaires/instruments we cannot rule out information bias. The use of self-reported academic scores is another limitation and future studies should use their academic records. Cross-sectional design of our study is yet another limitation since associations presented lack temporality. Prospective studies are necessary to study the associations between occurrence stressors and incidence of stress. Though response rates (80%) were fairly good there could be some selection (non-response) bias. Though participation was anonymous, authors had the opportunity to meet the non-respondents and ask them reasons for not participating in the survey. The main reasons that surfaced were the sensitive and personal nature of the study as well as the length of questionnaire. The other reasons were absence from college due to sickness or travel on the day of the survey.

Conclusion

The students reported higher levels of stress. The most frequently occurring stressors among the students were

related to academic and psychosocial domains. The associations between stressed cases and female gender, occurrence of academic and psychosocial stressors needs to be further tested by prospective studies.

Acknowledgements

The authors would like to thank Professor Dr Malik Hussain Mubbashar, Vice Chancellor University of Health Sciences, Lahore for his kind support, Waqas Sami, statistician UHS for his statistical help and Ahsan Shah, student CMH LMC for proof reading the draft. In addition the authors thank all the students who participated in this study for their valuable time given for completing the questionnaires.
Sources of funding: None

Author details

¹3rd year Undergraduate Medical Student, CMH Lahore Medical College, University of Health Sciences, Abdur Rahman Road, Lahore Cantt, Pakistan. ²Department of Physiology, CMH Lahore Medical College, University of Health Sciences, Abdur Rahman Road, Lahore Cantt, Pakistan. ³Department of Community Medicine, Melaka Manipal Medical College, Malaysia.

Authors' contributions

MS conceived, designed and carried out the study and prepared the first draft of the manuscript. SH helped design and carrying out the study. SM helped in carrying out the study, performed data analysis, and helped in drafting the manuscript. CTS assisted in design, interpretation of results, drafting the manuscript and critically evaluated earlier drafts of the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 2 April 2009

Accepted: 15 January 2010 Published: 15 January 2010

References

1. Dahlin M, Joneborg N, Runeson B: **Stress and depression among medical students: a cross sectional study.** *Med Educ* 2005, **39**:594-604.
2. Zocolillo M, Murphy GE, Wetzel RD: **Depression among medical students.** *J Affect Disord* 1986, **11**:91-96.
3. Tyssen R, Vaglum P, Gronvold NT, Ekeberg O: **Suicide ideation among medical students and youth physicians: a nationwide and prospective of prevalence and predictors.** *J Affect Disord* 2001, **64**:69-79.
4. Tyssen R, Hem E, Vaglum P, Gronvold NT, Ekeberg O: **The process of suicidal planning among medical doctors: predictors in a longitudinal Norwegian sample.** *J Affect Disord* 2004, **80**:191-198.
5. Malathi A, Damodaran A: **Stress due to exams in medical students-role of yoga.** *Indian J Physiol Pharmacol* 1999, **43**:218-24.
6. Bramness JA, Fixdal TC, Vaglum P: **Effect of medical school stress on the mental health of medical students in early and late clinical curriculum.** *Acta Psychiastr Scand* 1991, **84**:340-5.
7. Vitaliano PP, Russo J, Carr JE, Heerwagen JH: **Medical-school pressures and their relationship to anxiety.** *J Nerve Ment Dis* 1984, **172**:730-736.
8. Tyssen R, Vaglum P, Gronvold NT, Ekeberg O: **Factors in medical school that predict postgraduate mental health problems in need of treatment: A nationwide and longitudinal study.** *Med Educ* 2001, **35**:110-120.
9. Stewart SM, Lam TH, Betson CL, Wong CM, Wong AMP: **A prospective analysis of stress and academic performance in the first 2 years of medical school.** *Med Educ* 1999, **33**:243-50.
10. Fish C, Nies MA: **Health promotion needs of students in a college environment.** *Public Health Nurs* 1996, **13**:104-11.
11. Chew-Graham CA, Rogers A, Yassin N: **'I wouldn't want it on my CV or their records': medical students' experiences of help-seeking for mental health problems.** *Med Educ* 2003, **37**:873-80.
12. Silver HK, Glick AD: **Medical student abuse. Incidence, severity, and significance.** *JAMA* 1990, **263**:527-32.
13. Niemi PM, Vainiomaki PT: **Medical students' academic distress, coping and achievement strategies during the pre-clinical years.** *Teach Learn Med* 1999, **11**:125-34.

14. Vitaliano PP, Russo J, Carr JE, Heerwagen JH: **Medical school pressures and their relationship to anxiety.** *J Nerv Ment Dis* 1984, **172**:730-6.
15. Supe AN: **A study of stress in medical students at Seth G.S. Medical College.** *J Postgrad Med* 1998, **44**:1-6.
16. Saipanish R: **Stress among medical students in a Thai medical school.** *Med Teach* 2003, **25**:502-6.
17. Sherina MS, Rampal L, Kanesson N: **Psychological stress among undergraduate medical students.** *Med J Malaysia* 2004, **59**:207-11.
18. Shaikh BT, Kahloon A, Kazmi M, Khalid H, Nawaz K, Khan N, Khan S: **Students, stress and coping strategies: a case of Pakistani medical school.** *Educ Health (Abingdon)* 2004, **17**:346-53.
19. Cohen S, Kamarck T, Mermelstein R: **A global measure of perceived stress.** *J Health Soc Behav* 1983, **24**:385-96.
20. Hatcher L, Prus JS: **A measure of academic educational constraints: Out-of-Class circumstances that inhibit college student development.** *Educational and Psychological measurement* 1991, **51**:953-963.
21. Hammer LG, Grigsby TL, Woods S: **the conflicting demands of work, family and school among students of an urban university.** *The Journal of Psychology* 1998, **132**:220-227.
22. Devine TG: **Studying: skills, strategies, and systems.** *Handbook of Research on Teaching the English Language Arts.* New York Flood J, Jensen JM, Lapp D, Squire JR 1991, 743-53.
23. Amr M, Gilany AH, El-Hawary A: **Does gender predict students' stress in Mansoura, Egypt?.** *Med Educ Online* 2008, **13**:12.
24. Sreeramareddy CT, Shankar PR, Binu VS, Mukhopadhyay C, Ray B, Menezes RG: **Psychological morbidity, sources of stress and coping strategies among undergraduate medical students of Nepal.** *BMC Med Educ* 2007, **7**:26.
25. Ortmeier BG, Wolfgang AP, Martin BC: **Career commitment, career plans, and perceived stress: a survey of pharmacy students.** *Am J Pharm Educ* 1991, **55**:138-42.
26. Wolfgang AP: **The health professions stress inventory.** *Psychol Rep* 1988, **62**:220-2.
27. Wolfgang AP, Perri Wolfgang CF: **Job-related stress experienced by hospital pharmacists and nurses.** *Am J Hosp Pharm* 1988, **45**:1342-5.
28. Barnett CW, Hopkins WA, Jackson RA: **Burnout experienced by recent pharmacy graduates of Mercer University.** *Am J Hosp Pharm* 1986, **43**:2780-4.
29. Lapane KL, Hughes CM: **Job satisfaction and stress among pharmacists in the long-term care sector.** *Consult Pharm* 2006, **21**:287-92.
30. Wolf TM, Von Almen TK, Faucett JM, Randall HM, Franklin FA: **Psychosocial changes during first year of medical school.** *Med Educ* 1991, **25**:174-81.
31. Cohen S, Williamson G: **Perceived stress in a probability sample of the United States.** *The Social Psychology of Health: Claremont Symposium on Applied Social Psychology* Newbury Park, CA: SageSpacapam S, Oskamp S 1988, 31-67.
32. Chang EC: **Does dispositional optimism moderate the relationship between perceived stress and psychological well-being? A preliminary investigation.** *Personality and Individual Differences* 1998, **25**:233-40.
33. Hall NC, Chipperfield JG, Perry RP, Ruthig JC, Goetz T: **Primary and secondary control in academic development: gender-specific implications for stress and health in college students.** *Anxiety Stress Coping* 2006, **19**:189-210.
34. Otto MW, Fava M, Penava SJ, Bless E: **Life even, mood, and cognitive predictors of perceived stress before and after treatment for major depression.** *Cogn Ther Res* 1997, **21**:409-20.
35. Treadgold R: **Transcendent vocations: their relationship to stress, depression, and clarity of concept.** *J Humanistic Psychology* 1999, **39**:81-105.
36. Van Eck MM, Nicholson NA: **Perceived stress and salivary cortisol in daily life.** *Ann Behav Med* 1994, **16**:221-7.
37. Ebrecht M, Hextall J, Kirtley LG, Taylor A, Dyson M, Weinman J: **Perceived stress and cortisol levels predict speed of wound healing in health male adults.** *Psychoneuroendocrinology* 2004, **29**:798-809.
38. Cohen S, Tyrrell DA, Smith AP: **Negative life events, perceived stress, negative affect, and susceptibility to the common cold.** *J Pers Soc Psychol* 1993, **64**:131-40.
39. Gibson DM, Myers JE: **Perceived stress, wellness, and mattering: a profile of first-year citadel cadets.** *J College Student Development* 2006, **47**:647-60.
40. Matheny KB, Curlette WL, Aysan F, Herrington A, et al: **Coping resources, perceived stress, and life satisfaction among Turkish and American university students.** *Int J Stress Management* 2002, **9**:81-97.
41. Morrison R, O'Conner RC: **Predicting psychological distress in college students: the role of rumination and stress.** *J Clin Psychol* 2005, **61**:447-60.
42. Cohen S: **Laboratory for the Study of Stress, Immunity and Disease.** <http://www.psy.cmu.edu:16080/~scohen>.
43. Linn BS, Zeppa R: **Stress in junior medical students: relationship to personality and performance.** *J Med Educ* 1984, **59**:7-12.
44. Guthrie EA, Black D, Shaw CM, Hamilton J, Creed FH, Tomenson B: **Embarking upon a medical career: psychological morbidity in first year medical students.** *Med Educ* 1995, **29**:337-41.
45. Stewart SM, Betson C, Marshall I, Wong CM, Lee PWH, Lam TH: **Stress and vulnerability in medical students.** *Med Educ* 1995, **29**:119-27.
46. Ko SM, Kua EH, Fones CS: **Stress and the undergraduates.** *Singapore Med J* 1999, **40**:627-30.
47. Coles C: **Medicine and stress.** *Med Educ* 1994, **28**:3-4.
48. Redwood SK, Pollak MH: **Student-led stress management program for first-year medical students.** *Teach Learn Med* 2007, **19**:42-6.

Pre-publication history

The pre-publication history for this paper can be accessed here:<http://www.biomedcentral.com/1472-6920/10/2/prepub>

doi:10.1186/1472-6920-10-2

Cite this article as: Shah et al.: Perceived Stress, Sources and Severity of Stress among medical undergraduates in a Pakistani Medical School. *BMC Medical Education* 2010 **10**:2.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

